



Legionellosis

- Reported cases of disease, 1999-2003
- 5-year rate of disease, 1999-2003
- Reported cases by age group, 2001-2003
- Reported cases by gender, 2001-2003
- Reported cases by county, 2001-2003
- Reported cases by month, 2001-2003
- Reported cases by serogroup, 2001-2003

**Reported Cases of Legionellosis,
Rhode Island, 1999-2003**



**Legionellosis Rates
Rhode Island, 1999-2003**




Rates are based on 2000 Census

Note: Since the number of cases of meningococcal disease is less than 50 for one year, information has been combined for three years to allow demonstration of trends for the following figures:

Reported cases of Legionellosis by age group, Rhode Island, 2001-2003			
Age Group	Number of reported cases	Percent (%)	Rate ¹ of disease per 100,000
< 1	0	--	--
1-4	0	--	--
5-9	0	--	--
10-19	0	--	--
20-29	1	2.3	0.2
30-39	3	6.8	0.6
40-49	8	18.2	1.7
50-59	4	9.1	1.1
60-74	13	29.5	3.8
75+	15	34.1	6.4
TOTAL	44	100	1.4
¹ Denominators for computing rates for RI were based on 2000 Census counts obtained from the U.S. Census Bureau.			

Reported cases of Legionellosis, by gender, Rhode Island, 2001-2003			
Gender	Number of reported cases	Percent (%)	Rate ¹ of disease per 100,000
Males	27	61.4	1.8
Females	17	38.6	0.9
TOTAL	44	100	1.4
¹ Denominators for computing rates for RI were based on 2000 Census counts obtained from the U.S. Census Bureau.			

Reported cases of Legionellosis by County, Rhode Island, 2001-2003			
County	Number of reported cases	Percent (%)	Rate ¹ of disease per 100,000
Bristol	1	2.3	0.7
Kent	10	22.7	2.0
Newport	4	9.1	1.6
Providence	27	61.4	1.4
Washington	2	4.5	0.5
TOTAL	44	100	1.4
¹ Denominators for computing rates for RI were based on 2000 population counts obtained from the U.S. Census Bureau.			



Questions regarding these data can be directed to the
Office of Communicable Diseases at (401) 222-2577